

On Supplements in Modules

by

Jutta HAUSEN and Johnny A. JOHNSON¹

(Received March 6, 1981)

0. Introduction

In a recent paper, K. Varadarajan established a criterion under which supplements in projective modules are direct summands. The purpose of this note is to weaken Varadarajan's condition, and to extend the result to quasi-projective modules. Our criterion will be the Hopfity of the radical. Hopfian quasiprojective modules will be characterized and criteria relating the Hopfity of a module to that of its radical will be given. In addition, we will indicate alternate short proofs of Varadarajan's characterization of perfect and semi-perfect rings.

1. Supplements with Hopfian radical

Notation and terminology will follow Varadarajan [9] and Anderson-Fuller [1]. All modules will be unital left modules. A module M is said to be *Hopfian* if every surjective endomorphism of M is an automorphism.

LEMMA 1.1. *Let $f: M \rightarrow M$ be an epimorphism and let $\text{Ker } f \subseteq J(M)$. If $J(M)$ is Hopfian then f is an automorphism.*

Proof. By ([1]; p. 121, 9.15]), $J(M) = f(J(M))$. Since $J(M)$ is Hopfian, $0 = \text{Ker } f \cap J(M) = \text{Ker } f$.

COROLLARY 1.2. *Suppose $J(M)$ is Hopfian. Then M is Hopfian if and only if every submodule N of M with $M/N \simeq M$ is small in M .*

Proof ([1]; p. 120, 9.13). Let N and H be submodules of M , and let $M = N + H$. If N and H are supplements of each other and M is quasi-projective, then $M = N \oplus H$ ([7]; p. 92, 2.3). However, even in a free module a given submodule need not have a supplement ([4]; Theorem 4.17); also, if H is a supplement, the existence of N so that H and N are mutual supplements is not certain; in particular, we do not know whether a supplement in a quasi-projective module is quasi-projective.

¹ The second author was partially supported by a University of Houston Central Campus Research Enabling Grant.

Classification (1978 Math Reviews Index): 13C10, 13C05, 13C13

The following result contains a criterion under which supplements in quasi-projective modules are direct summands and, thus, are quasi-projective ([11]; p. 442, 2.3).

THEOREM 1.3. *Let Q be a quasi-projective module, and let $H \subseteq Q$ be a supplement of some submodule of Q . If $J(H)$ is Hopfian then H is a direct summand of Q ; in particular, H is quasi-projective.*

Remark. It is well known that every Noetherian module is Hopfian; also, finitely generated modules over commutative rings are Hopfian ([10]; p. 506, 1.2). Thus, Theorem 1.3 generalizes ([9]; p. 563, 2.4). That Hopfian modules in turn need not be finitely generated can be seen from the infinite Hopfian p -primary group constructed by R. S. Pierce ([8]; p. 302, 16.4).

Proof of Theorem 1.3. Let H be a supplement of $N \subseteq Q$. Then $Q = N + H$ and $N \cap H$ is small in H ([7]; p. 87, 1.3). By ([7]; p. 91, 2.1) there exists a homomorphism $\psi: Q \rightarrow H$ making the diagram

$$\begin{array}{ccc} & Q & \\ & \downarrow \psi & \\ H & \xrightarrow{\alpha} & Q/N \end{array}$$

commutative, where v is the natural epimorphism and $\alpha = v|_H$. From $Q/N = v(H) = \alpha(\psi(H))$, it follows that $H = \psi(H) + H \cap N$, so that $H = \psi(H)$. Let $f = \psi|_H$. Then $\text{Ker } f \subseteq H \cap N \subseteq J(H)$, and f is an automorphism by Lemma 1.1. Since $\psi \circ f^{-1} = 1_H$, H is a direct summand.

2. Quasi-projective Hopfian modules

Theorem 1.3 motivates the consideration of quasi-projective modules with Hopfian radical. We have the following easy characterization of quasi-projective Hopfian modules.

PROPOSITION 2.1. *The quasi-projective module Q is Hopfian if and only if Q is not isomorphic to a proper direct summand of Q .*

Proof ([3]; p. 6, Lemma 4). If R is a ring and $J(R) = 0$ then $J(P) = 0$ for every projective R -module P ([1]; p. 196, 17.10). Thus $J(P)$ being Hopfian does not imply the Hopficity of P . At the other end of the scale, however, the situation is different:

PROPOSITION 2.2. *Let Q be a quasi-projective module such that $J(S) \neq 0$ for every non-zero direct summand S of Q . Then $J(Q)$ Hopfian implies Q Hopfian.*

Proof. Suppose $Q = C \oplus S$, $C \simeq Q$. Then $J(Q) = J(C) \oplus J(S)$ ([1]; p. 121, 9.19), and $J(C) \simeq J(Q)$. Since $J(Q)$ is Hopfian, $J(S) = 0$, so that $S = 0$.

COROLLARY 2.3. *Let Q be a quasi-projective module such that $J(S) \neq 0$ for every direct summand $S \neq 0$ of Q , and let $H \subseteq Q$ be a supplement in Q . If $J(H)$ is Hopfian, then H is a quasi-projective Hopfian direct summand of Q .*

Proof. Theorem 1.3 and Proposition 2.2.

3. Pefect and semi-perfect rings

The ring R is called left (semi-) perfect if every (finitely generated) left R -module has a projective cover ([2]; p. 471). While left semi-perfect rings are right semi-perfect, left perfect rings need not be right perfect.

For P a projective module, the following conditions are known to be equivalent ([6]; p. 526, Satz; and [7]; p. 96, 3.3): (i) P has property (P_1) ; (ii) every epimorphic image of P has a projective cover; (iii) P has property (P_2) . Properties (P_1) and (P_2) are inherited by epimorphic images ([5]; p. 245, 11.1.4 (3); and [7]; p. 90, 1.12), and sums of finitely many modules satisfying (P_1) satisfy (P_1) ([5]; p. 268, Exercise 4b).

These remarks immediately prove Varadarajan's Theorems 1.6 and 1.7 ([9]; p. 561).

References

- [1] ANDERSON, F. W. and FULLER, K. R.; "Rings and Categories of Modules," Springer-Verlag, New York, 1974.
- [2] BASS, H.; Finitistic dimension and a homological generalization of semi-primary rings, *Trans. Amer. Math. Soc.*, **95** (1960), 466–488.
- [3] FUCHS, L. and RANGASWAMY, K. M.; Quasi-projective abelian groups, *Bull. Soc. Math. France*, **98** (1970), 5–8.
- [4] HAUSEN, J.; Supplemented modules over Dedekind domains, *Pacific J. Math.*
- [5] KASCH, F.; "Moduln und Ringe," B. G. Teubner, Stuttgart, 1977.
- [6] KASCH, F. and MARES, E.; Eine Kennzeichnung semi-perfekter Moduln, *Nagoya Math. J.*, **27** (1966), 525–529.
- [7] MIYASHITA, Y.; Quasi-projective modules, perfect modules, and a theorem for modular lattices, *J. Fac. Sci. Hokkaido Univ.*, **19** (1966), 86–110.
- [8] PIERCE, R. S.; Homomorphisms of primary abelian groups, *Topics in Abelian Groups*, Chicago, 1963, pp. 215–310.
- [9] VARADARAJAN, K.; Modules with supplements, *Pacific J. Math.*, **82** (1979), 559–564.
- [10] VASCONCELOS, W. V.; On finitely generated flat modules, *Trans. Amer. Math. Soc.*, **138** (1969), 505–512.
- [11] WU, L. E. T. and JANS, J. P.; On quasi projectives, *Illinois J. Math.*, **11** (1967), 439–448.

Department of Mathematics
University of Houston, Central Campus
Houston, Texas 77004
U.S.A.